



www.AmericanHeadacheSociety.org

Patient Education Kit for Migraine

AHS Complementary and Integrative Medicine Special Interest Section

Exercise

Regular exercise reduces headache frequency, severity, and duration.^{1,2,3,4} There are positive effects on neuroinflammatory, neurovascular, and psychological systems during exercise, which contribute to improvements in headache.⁵ Exercise is beneficial when performed on its own, however it may have even more impact when used in combination with other preventive treatments, such as medication.^{6,7} Evidence-based suggestions to consider are as follows:


- **Sustainable exercise:** Be kind to yourself. Develop an exercise program that is sustainable, enjoyable, and convenient. A combination of aerobic exercise (e.g., brisk walking, running, swimming) and resistance training (e.g., yoga, Pilates, weight-lifting) is recommended.^{4,8,9} Choose exercises you can tolerate, such as low-impact exercise that are gentle on your body. Vary your exercise routine to keep it fun; this can help you stick with a regular exercise plan over time.¹⁰ Be patient and stick with it. Most programs typically take 8 weeks to impact headaches positively.^{11,12}
- **Exercise Parameters:** Start with slow, gentle exercise, and increase the intensity, frequency, and duration of exercise over time as endurance improves. Higher-intensity training has increased benefit for headache prevention.³ Strive to exercise for 30 minutes per day, at least 3-5 days per week.^{8,9} Include muscle-strengthening activities at least two days per week using your own body weight, free weights, resistance machines or bands to train the entire body.^{8,9,11}


Methods for Determining Exercise Parameters:

Exercise Parameters	Light	Moderate	Vigorous
Duration ^{8,9,11}	Below symptoms onset	150-300 minutes a week	75-150 minutes a week
Frequency ^{8,9,11,12}	3 or more days / week	3 or more days / week	3 or more days / week
Intensity (based upon Talk Test parameters) ^{13,14}	Can talk easily	Can talk, not sing	Difficulty maintaining conversation

- **Food and Liquid Intake:** Maintain good hydration before, during, and after exercise. Even mild dehydration can trigger headaches.¹⁵ Eat a balanced snack (including protein, fat, and carbohydrates) before exercise.
- **Neck pain:** Many individuals with headache also have neck pain. Aerobic exercise has been shown to reduce neck pain, as well as migraine and tension-type headache.¹⁶ Be mindful of the positioning of your head and neck during exercise. If you need additional support with neck pain, speak with your physician.
- **Postural Recommendations:** Below are simple recommendations to consider during daily activities to improve your posture and assist with headache management ^{17,18,19}

- Switch positions frequently by scheduling timed breaks. Stand up to reset your body
- Consider adjusting your workstation by modifying computer equipment and furniture when possible.
 - When using a laptop, position the monitor at eye level. An external keyboard and mouse can help provide appropriate set up.
 - When using books or smaller electronic devices bring the object to eye level.
- Position frequently used objects within easy reach to limit repetitive stretching or twisting.





- **Exercises for Postural Reset:** Complete exercise throughout the day to improve upright posture and self-awareness.^{4,11,20, 21}

- While sitting or standing, bend both elbows to 90 degrees with palms facing up.
- Relax your shoulders and gently move your hands out to the sides while keeping your elbows close to your body. Only move within your available/comfortable range.
- Return to the starting position.
- This can be done a few times a day.
- This should not be irritating or heightening your headache, discontinue if that occurs.



- **Relaxation Training:** The practice of yoga has been shown to decrease headache frequency, intensity, disability, and use of rescue medications, when used in combination with standard migraine treatments, such as medications and lifestyle changes. The practice of yoga, when practiced 60 minutes per day, on 3-5 days per week, has been shown to decrease headache frequency.²² Yoga can be performed in the home or in a teaching center.

References

1. Lemmens J, De Pauw J, Van Soom T, Michiels S, Versijpt J, Van Breda E, Castien R, De Hertogh W. The effect of aerobic exercise on the number of migraine days, duration and pain intensity in migraine: a systematic literature review and meta-analysis. *The journal of headache and pain*. 2019 Dec;20(1):1-9.
2. Machado-Oliveira L, da Silva Gauto YO, de Santana Neto FJ, da Silva MG, Germano-Soares AH, Diniz PR. Effects of different exercise intensities on headache: a systematic review. *American Journal of Physical Medicine & Rehabilitation*. 2020 May 1;99(5):390-6.
3. Barber M, Pace A. Exercise and migraine prevention: a review of the literature. *Current Pain and Headache Reports*. 2020 Aug;24(8):1-7.
4. Tsou AY, Datko M, Oristaglio J, Patient MP. Nonpharmacologic Interventions for Migraine Prevention.
5. Irby MB, Bond DS, Lipton RB, Nicklas B, Houle TT, Penzien DB. Aerobic exercise for reducing migraine burden: mechanisms, markers, and models of change processes. *Headache: The Journal of Head and Face Pain*. 2016 Feb;56(2):357-69.
6. Varkey E, Cider Å, Carlsson J, Linde M. Exercise as migraine prophylaxis: a randomized study using relaxation and topiramate as controls. *Cephalalgia*. 2011 Oct;31(14):1428-38.
7. Santiago MD, Carvalho DD, Gabbai AA, Pinto MM, Moutran AR, Villa TR. Amitriptyline and aerobic exercise or amitriptyline alone in the treatment of chronic migraine: a randomized comparative study. *Arquivos de neuro-psiquiatria*. 2014;72:851-5.
8. Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, King AC, Macera CA, Castaneda-Sceppa C. Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Circulation*. 2007;116(9):1094.
9. Garber, C.E., Blissmer, B., Deschenes, M.R., Franklin, B.A., Lamonte, M.J., Lee, I.M., Nieman, D.C. and Swain, D.P., 2011. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Medicine and science in sports and exercise*, 43(7), pp.1334-1359.
10. Lakicevic N, Gentile A, Mehrabi S, Cassar S, Parker K, Roklicer R, Bianco A, Drid P. Make fitness fun: could novelty be the key determinant for physical activity adherence?. *Frontiers in Psychology*. 2020:2558.
11. Sari Aslani P, Hassanpour M, Razi O, Knechtle B, Parnow A. Resistance training reduces pain indices and improves quality of life and body strength in women with migraine disorders. *Sport Sciences for Health*. 2021 Oct 4:1-1.
12. Patel P, Minen MT. Complementary and integrative health treatments for migraine. *Journal of neuro-ophthalmology: the official journal of the North American Neuro-Ophthalmology Society*. 2019 Sep;39(3):360.
13. Sørensen L, Larsen KS, Petersen AK. Validity of the talk test as a method to estimate ventilatory threshold and guide exercise intensity in cardiac patients. *Journal of Cardiopulmonary Rehabilitation and Prevention*. 2020 Sep 1;40(5):330-4.
14. Porcari JP, Falck-Wiese K, Suckow-Stenger S, Turek J, Wargowsky A, Cress ML, Doberstein ST, Hahn CJ, Ryskey AL, Foster C. Comparison of the talk test and percent heart rate reserve for exercise prescription. *Kinesiology*. 2018 Jun 21;50(1):3-10.

15. Jones CW, Remboski LB, Freeze B, Braz VA, Gaughan JP, McLean SA. Intravenous fluid for the treatment of emergency department patients with migraine headache: a randomized controlled trial. *Annals of Emergency Medicine*. 2019 Feb 1;73(2):150-6.
16. Krøll LS, Hammarlund CS, Linde M, Gard G, Jensen RH. The effects of aerobic exercise for persons with migraine and co-existing tension-type headache and neck pain. A randomized, controlled, clinical trial. *Cephalalgia*. 2018 Oct;38(12):1805-16.
17. Heidarimoghadam R, Mohammadfam I, Babamiri M, Soltanian AR, Khotanlou H, Sohrabi MS. What do the different ergonomic interventions accomplish in the workplace? A systematic review. *International Journal of Occupational Safety and Ergonomics*. 2022 Jan 2;28(1):600-24.
18. Park SY, Yoo WG. Effects of the sustained computer work on upper cervical flexion motion. *Journal of physical therapy science*. 2014;26(3):441-2.
19. Smith L, Louw Q, Crous L, Grimmer-Somers K. Prevalence of neck pain and headaches: impact of computer use and other associative factors. *Cephalalgia*. 2009 Feb;29(2):250-7.
20. Florencio LL, de Oliveira IV, Lodovichi SS, Bragatto MM, Benatto MT, Dach F, Fernández-de-Las-Peñas C, Bevilaqua-Grossi D. Cervical muscular endurance performance in women with and without migraine. *Journal of orthopaedic & sports physical therapy*. 2019 May;49(5):330-6.
21. Ashina S, Bendtsen L, Lyngberg AC, Lipton RB, Hajiyeva N, Jensen R. Prevalence of neck pain in migraine and tension-type headache: a population study. *Cephalalgia*. 2015 Mar;35(3):211-9.
22. Kumar A, Bhatia R, Sharma G, Dhanlika D, Vishnubhatla S, Singh RK, Dash D, Tripathi M, Srivastava MVP. Effect of yoga as add-on therapy in migraine (CONTAIN): A randomized clinical trial. *Neurology*. 2020 May 26;94(21):e2203-e2212.